


Additional information and updates can be found online at http://go.usa.gov/xkMBc



If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)

1969 - Landfill begins operation

- Former Industrial Waste Disposal Area (1969 to early 1980s)
 - Disposed liquid wastes included solvents, oily sludges, and thinners
- Navy Municipal Landfill Area (1969 to 1992)

1990 - Landfill designated as Superfund site

Chlorinated volatile organic compounds, including vinyl chloride, present in groundwater

1993 - Landfill clean-up begins

- Engineered landfill cap constructed
- Monitoring and operation of a groundwater treatment system to address chlorinated volatile organic compounds (including breakdown products such as vinyl chloride)
- Ongoing monitoring activities have also included sampling offsite private wells and adding offsite monitoring wells to evaluate the migration of contaminants (including vinyl chloride and trichloroethene) in groundwater from Area 6

2001 – Excavation at Former Industrial Waste Disposal Area

Approximately 3,472 tons of impacted soil excavated and properly disposed

2010 – Washington State begins regulation of 1,4-dioxane

- 1,4-Dioxane present in groundwater onsite and offsite
- 1,4-Dioxane not part of original clean-up effort in 1993
 - 1,4-Dioxane historically added to certain chlorinated solvents as a stabilizer
 - Existing groundwater treatment system does not remove 1,4-dioxane
- Refining and finalizing clean-up strategy

2017 - PFAS found in groundwater at Area 6

2018 – Additional Investigation – Area 6 Offsite Sampling

- To assess whether vinvl chloride and 1.4-dioxane in groundwater have migrated farther to the south and/or west
- To assess whether PFAS in groundwater at Area 6 has migrated offsite

LEGEND

- Area 6 boundary
- Base boundary
 - PFOS perfluorooctane sulfonate ppb parts per billion
- parts per trillion ppt

aqueous film forming foam EPA U.S. Environmental Protection Agency

AFFF

PFAS per- and polyfluoroalkyl substances perfluorooctanoic acid


PFOA





Why Is the Navy Sampling for Vinyl Chloride at Area 6?

Additional information and updates can be found online at http://go.usa.gov/xkMBc If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)



The Navy is being protective by identifying potential vinyl chloride in drinking water. The Navy is taking action for past disposal practices.

- Monitoring indicates that the vinyl chloride groundwater plume extends off-base.
- The EPA has established a maximum contaminant level for drinking water of 2 ppb for vinyl chloride.
- The Navy will sample off-base drinking water to evaluate potential human health risk associated with vinyl chloride from Area 6 and will take action as necessary.


LEGEN	D	Notes:		
Vinyl chloride groundwater plume (≥ 0.10 ppb)		 Results in exceedance of the federal maximum contaminant level (2 ppb) are shown in red. 		
— Area 6 boundary				
🗔 Base boundary		2. Data shown in figure is from 2017.		
Well		NA = well not analyzed for vinyl chloride		
 Approximate groundwater flow direction 		ND = vinyl chloride not detected		
AFFF	aqueous film forming foam	PFOS	perfluorooctane sulfonate	
epa Pfas Pfoa	U.S. Environmental Protection Agency per- and polyfluoroalkyl substances perfluorooctanoic acid	ppb ppt	parts per billion parts per trillion	





Why Is the Navy Sampling for 1,4-Dioxane at Area 6?

Additional information and updates can be found online at http://go.usa.gov/xkMBc If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)



The Navy is being protective by identifying potential 1,4-dioxane in drinking water. The Navy is taking action for past disposal practices.

- Monitoring indicates that the 1,4-dioxane groundwater plume extends off-base.
- The EPA has not established a maximum contaminant level for 1,4-dioxane. The Navy is adopting 35 ppb as the drinking water exposure assessment action level.
- The Navy will sample off-base drinking water to evaluate potential human health risk associated with 1,4-dioxane from Area 6 and will take action as necessary.

LEGEND

- 1,4-Dioxane groundwater plume (≥ 0.44 ppb)
- Area 6 boundary
- Base boundary
- Well
- Approximate groundwater flow direction

Notes:

- 1. There were no results in exceedance of the EPA value for cancer risk in tap water (35 ppb).
- 2. Data shown in figure is from 2017.
- NA = well not analyzed for 1,4-dioxane
- ND = 1,4-dioxane not detected

AFFF aqueous film forming foam

- EPA U.S. Environmental Protection Agency
- PFAS per- and polyfluoroalkyl substances
- PFOA perfluorooctanoic acid

- PFOS perfluorooctane sulfonate
- ppb parts per billion
- ppt parts per trillion





Why Is the Navy Sampling for PFAS Off-Base at Area 6?

Additional information and updates can be found online at http://go.usa.gov/xkMBc If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)



The Navy is being protective by identifying potential unregulated compounds (PFAS) in drinking water. The Navy is taking action for past disposal practices.

- The Navy used AFFF, a source of PFAS, for fire fighting.
- The EPA established a lifetime health advisory (70 ppt) for two PFAS, specifically PFOS and PFOA, in drinking water.
- Navy policy is to identify and prioritize locations with the potential for exposure to PFOS and/or PFOA.
- PFOS and PFOA have been identified in groundwater at Area 6.
- The Navy will sample off-base drinking water to evaluate potential human health risk associated with PFOS and PFOA from Area 6 and will take action as necessary.

	ID rea 6 boundary ase boundary	Notes: 1. Results in exceedance of the EPA lifetime health advisory for PFOS and/or PFOA (70 ppt) are shown in red.		
 Well Approximate groundwater flow direction 		 2. Data shown in figure is from 2017. ND = not detected ND = PFOS concentration (top) 1.82 ppt = PFOA concentration (bottom) 		
AFFF EPA PFAS PFOA	aqueous film forming foam U.S. Environmental Protection Agency per- and polyfluoroalkyl substances perfluorooctanoic acid	PFOS ppb ppt	perfluorooctane sulfonate parts per billion parts per trillion	





Additional information and updates can be found online at http://go.usa.gov/xkMBc If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)



LEGEND

Base boundary Area 6 boundary Area 6 sampling area boundary
 Area 6 off-base sampling area parcel

Area 6 Off-Base Decision-Making Process


- The Off-Base Sampling Area includes parcels within 1/2 mile in the direction of groundwater flow to the south and parcels within 1/2 mile of the western landfill boundary.
- If your property is in the sampling area and you have a water well, then the Navy will sample your well for PFOS, PFOA, 1,4-dioxane, and vinyl chloride.
- The Navy will provide an alternate water source if your drinking water well exceeds the action levels for PFOS, PFOA, 1,4-dioxane, and/or vinyl chloride.
- Step-out sampling may be conducted based on the results.

AFFF aqueous film forming foam

- EPA U.S. Environmental Protection Agency
- PFAS per- and polyfluoroalkyl substances

PFOA perfluorooctanoic acid

- PFOS perfluorooctane sulfonate
- ppb parts per billion
- ppt parts per trillion








Vinyl Chloride and 1,4-Dioxane Plumes at Area 6

Additional information and updates can be found online at http://go.usa.gov/xkMBc

If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)





Data shown in figures is from 2017. ND = not detected

ppt

parts per trillion





SEPA Wingen Sur Dynamid Wingen S

Additional information and updates can be found online at http://go.usa.gov/xkMBc

Where Does Vinyl Chloride Come From?

- It is a manufactured chlorinated solvent with no natural occurrence.
- It is formed when larger chlorinated solvents, such as trichloroethene (TCE) or tetrachloroethene (PCE) break down in the environment. TCE, PCE, and other chlorinated solvents were used to clean aircraft and parts.
- It is mostly used in the production of polyvinyl chloride (PVC) to make plastic, pipes, wire coatings, and packaging materials.
- Smaller amounts are used in furniture and automobile upholstery, wall coverings, housewares, and automotive parts; it was also used in the past as a refrigerant.
- Low levels are found in tobacco smoke.

Sources and Potential Exposure

- For most people, exposure occurs when they:
 - Drink contaminated groundwater. Chlorinated solvents disposed of or spilled on the ground can migrate into the groundwater.
 - Eat food in contact with contaminated packaging; this exposure route has decreased.
 - Breathe air after evaporation from contaminated water used for bathing or laundering.
 - Breathe exhaust from factories or evaporation from storage areas.
 - Come into skin contact with contaminated groundwater or products containing vinyl chloride; this exposure route is not significant.
- People who work with vinyl chloride and other solvents may have the highest exposures from inhaling vapors.
- Vinyl chloride is eliminated quickly from the body and does not accumulate in plants or animals. It has not been found in breast milk.

If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)

Health Effects

- The health effects in people drinking water or breathing air contaminated with low levels of vinvl chloride are not well understood. Animal studies report:
 - Effects on liver, kidney, and nervous system
 - Possible effects on fetal growth and development; birth defects not seen
 - Possible damage to sperm and testes
- Vinyl chloride is a known liver carcinogen. Animal studies suggest infants and young children may be more susceptible than adults to cancer induced by vinyl chloride exposure.

Vinyl Chloride Action Level in Water

- EPA has set an enforceable maximum contaminant level in drinking water of 2 ppb.
- Maximum contaminant levels are set as close to health goals as possible but consider cost to remove and ability to detect.
- The Navy will provide bottled water if drinking water has vinyl chloride levels higher than 2 ppb.
 - Non-cancer effects at this level are not expected.
 - Drinking at this level over a lifetime adds cancer risk of one additional cancer in 10,000 people.

- AFFF aqueous film forming foam
- ATSDR Agency for Toxic Substances and Disease Registry PFOS EPA
- U.S. Environmental Protection Agency PFAS per- and polyfluoroalkyl substances
- perfluorooctane sulfonate

PFOA

parts per billion ppb

perfluorooctanoic acid

ppt parts per trillion



Additional information and updates can be found online at http://go.usa.gov/xkMBc

If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)

Where Does 1,4-Dioxane Come From?

- It is used in the production of solvents and plastics.
 - In the past, it was added to stabilize chlorinated solvents, including those used to clean aircraft and parts.
 - It is a by-product in antifreeze and aircraft deicing fluids.
 - It is a by-product in clear palastic such as water or soda bottles.
- Very low levels are found in consumer products and food.
- Very low levels are found in cosmetics, detergents, and shampoos.
 - Very low levels are found in food supplements, packaging adhesives, or food crops treated with pesticides that contain 1,4-dioxane.
 - It is used in the manufacture of some pharmaceuticals.

Sources and Potential Exposure

- Most 1,4-dioxane contamination found in groundwater comes from leaking storage tanks, discharges, and past disposal practices.
- It does not break down easily in water and travels ahead of other contaminants in the groundwater.
- For most people, exposure occurs when they:
 - Drink contaminated groundwater.
 - Eat food in contact with contaminated packaging; this exposure route has decreased.
 - Breathe air after evaporation from contaminated water used for bathing or laundering.
 - Come into skin contact with contaminated groundwater or products containing 1,4-dioxane.
- 1,4-Dioxane is broken down and eliminated quickly from the body. It does not accumulate in plants or animals. It may be transferred to breast milk.

Health Effects

- The health effects in people drinking water contaminated with 1,4-dioxane are not well understood. Animal studies report:
 - Effects on liver and kidney
 - A slight effect on fetal growth and development; birth defects not seen
- People breathing low levels of 1-4, dioxane for short periods of time have reported eye and nose irritation.
- EPA has classified 1,4-dioxane as likely to be carcinogenic to humans by all exposure routes. Animals given 1,4-dioxane in drinking water have had increased numbers of tumors of the nasal cavity, liver, and gall bladder.

1,4-Dioxane Action Level in Water

- EPA has not set an enforceable maximum contaminant level for 1.4-dioxane.
- EPA has set a non-enforceable lifetime health advisory of 200 ppb for 1.4-dioxane, which:
 - Protects the general population over a lifetime; non-cancer health effects are not expected.
 - Assumes 20% of exposure from water and 80% from other sources.
- The Navy will provide bottled water if drinking water has 1,4-dioxane levels higher than 35 ppb.
 - Non-cancer effects at this level are not expected.
 - Drinking at this level over a lifetime adds cancer risk of one additional cancer in 10,000 people.
- AFFF aqueous film forming foam

EPA

PFAS

- ATSDR Agency for Toxic Substances and Disease Registry PFOS

PFOA

- ppt parts per trillion
- U.S. Environmental Protection Agency per- and polyfluoroalkyl substances
- perfluorooctane sulfonate parts per billion ppb

perfluorooctanoic acid





Per- and Polyfluoroalkyl Substances (PFAS)

Additional information and updates can be found online at http://go.usa.gov/xkMBc

35 -

30

ŝ.,

a 25

20

If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)

Where Do PFAS Come From?

- They are manufactured compounds with no natural occurrence.
- They have been used since the 1950s in many products.
- They last a long time in the environment.
- They are globally distributed and have been detected in people, wildlife, and fish.



AFFF, paints and stains, water-repelling fabrics





nonstick cookware, food packaging, stain-resistant carpets

Sources and Potential Exposure

- Monitoring by the Centers for Disease Control estimates that most people in the U.S. have PFAS in their bodies.
- PFAS can build up in the body until exposure stops.
- For most people, exposure occurs when they:
 - Ingest contaminated food, water, or soil.
 - Breathe air that contains contaminated dust from carpets, upholstery, clothing, dirt, etc.
- PFAS reach the fetus or nursing infant of a mother who is exposed.
- Exposure is not significant through skin contact or inhalation when bathing or showering.

Levels of PFOS and PFOA in blood serum of the U.S. population \$ 30.2 since their phase-out from products began around 2002







Health Effects

- More research is needed to confirm or rule out possible links between exposure and health effects. Based on limited evidence from studies with people, the potential health effects include:
 - Increased cholesterol levels
 - Changes in growth, learning, and behavior of the developing fetus and child
- Decreased fertility
- Altered thyroid function
- Increased risk of certain types of cancer
- Changes in the immune system
- Animals in PFAS studies exhibit developmental, reproductive, and liver effects, as well as an increased occurrence of cancer.
- The presence of PFAS in a well or water system does not predict whether any health effects will occur.

PFAS Action Level in Water

- EPA has not set an enforceable maximum contaminant level for any PFAS.
- EPA has set a lifetime health advisory of 70 ppt in drinking water for PFOS and PFOA combined, which:
 - Protects sensitive populations including the fetus or nursing infant of a mother who is exposed.
 - Protects the general population for a lifetime.
 - Assumes 20% of exposure from drinking water and 80% from other sources.
 - Is based on animal studies and considers health effects found in exposed populations.
- The Navy will provide bottled water if drinking water has PFOS or PFOA levels higher than 70 ppt individually or combined.
- AFFF aqueous film forming foam

PFAS

- Agency for Toxic Substances and Disease Registry PFOS ATSDR EPA
 - U.S. Environmental Protection Agency per- and polyfluoroalkyl substances
 - ppb

PFOA

- ppt
- parts per billion parts per trillion

perfluorooctanoic acid

perfluorooctane sulfonate






SEPA Waingen Size Dynamical of Next Steps

Additional information and updates can be found online at http://go.usa.gov/xkMBc

If you have specific questions, please contact PAO feedback@navv.mil (email) or 360-396-1030 (voicemail)

Is the result above the action level?



Is the result at or below the action level?

No immediate action is needed: the Navy may monitor.

Please help us to:

- Identify water wells in sampling area.
- Sample wells to aid site investigations.
- Understand the community's concerns by participating in the Restoration Advisory Board.

Additional Follow-On Actions

- Meet with homeowners with results above action levels to gather data needed to develop long-term solutions.
- Continue to communicate with residents through the press, websites, emails, phone line, and Restoration Advisory Board.
- Continue to partner with local, state, and federal environmental and health agencies to determine the best path forward.
- Continue to investigate groundwater for PFAS, vinyl chloride, and 1,4-dioxane source locations, migration, and exposure pathways.
- Continue to monitor the science and regulations related to PFAS, vinyl chloride, and 1,4-dioxane.
- Refine and finalize clean-up strategy for 1,4-dioxane.

AFFF aqueous film forming foam FPA U.S. Environmental Protection Agency PFAS per- and polyfluoroalkyl substances PFOA perfluorooctanoic acid

perfluorooctane sulfonate PFOS ppb parts per billion

parts per trillion ppt





We Need Your Help – Drinking Water Sampling Process

Additional information and updates can be found online at http://go.usa.gov/xkMBc

If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)

Sampling Process

- We encourage you to contact us if you've received a sampling notification letter and have not signed up for sampling.
- We need your help to:
 - Make your appointment (sampling will take less than an hour).
 - Review and fill out the questionnaire.
- A team of qualified professionals will:
 - Collect cold water from the sample point (water will run for 3–5 minutes).
 - Analyze the sample according to EPA guidelines for a sampling and analysis process that follows strict quality control and quality assurance protocols.



Other Ways to Schedule an Appointment

To schedule an appointment for sampling your drinking water, please contact: PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)

AFFFaqueous film forming foamEPAU.S. Environmental Protection Agency

PFAS per- and polyfluoroalkyl substances PFOA perfluorooctanoic acid PFOS perfluorooctane sulfonate ppb parts per billion ppt parts per trillion



Sign Up for Your Sampling Appointment Here

Additional information and updates can be found online at http://go.usa.gov/xkMBc

If you have specific questions, please contact PAO_feedback@navy.mil (email) or 360-396-1030 (voicemail)

Off-Base Drinking Water Well Sampling

- Drinking water samples will be collected Monday–Friday, February 5–16.
- Sampling appointments are available 8 a.m.–7 p.m. Monday–Friday and 9 a.m.–1 p.m. Saturday–Sunday.
- The homeowner must give permission for sampling.
- Sampling takes less than an hour.
- An adult resident (18 years of age or older) must be present during sampling.

